

Detection of agricultural droughts in the Philippines within the 2015-2016 El Niño episode

Marco D. Macapagal
MS Environmental Science student
Institute of Environmental Science and Meteorology
University of the Philippines Diliman

Dr. Gay Jane P. Perez
Thesis Adviser
Institute of Environmental Science and Meteorology
College of Science
University of the Philippines Diliman

Dr. Felino P. Lansigan
Thesis Reader
College of Arts and Sciences
University of the Philippines Los Baños

ABSTRACT

Drought is a natural hazard characterized by below-normal precipitation, which can persist for months or even years. This recurrent extreme climate event can cause crop damage and yield loss, thereby inflicting negative socioeconomic impacts all over the world. As an agricultural country, the Philippines is highly susceptible to adverse impacts of drought events. Unfortunately, the country has no established method that can fully characterize agricultural drought. The Philippine Atmospheric, Geophysical and Astronomical Services Administration currently uses reliable indices intended for detecting and monitoring meteorological drought. In this study, the primary objective is to develop a simple, accurate, cost-effective, and reliable satellite-derived drought index capable of detecting agricultural drought events in the Philippines within the recent Strong El Niño episode. Specifically, it aims to determine the accuracy of this index in detecting the occurrence of agricultural drought events, evaluate the performance of this remotely sensed agricultural drought index by comparing it with rainfall-based drought indices, and compare the trends of rice and corn crop yield anomalies with the proposed drought index. This paper also tries to determine the implications of having an agricultural drought index on drought assessment, decision-making and management. Land products acquired from the Terra MODIS, such as Land Surface Temperature and Normalized Difference Vegetation Index, were employed to devise a remotely sensed localized agricultural drought index called Standardized Vegetation and Temperature Ratio (SVTR). Monthly SVTR maps were produced from January 2015 to December 2016. The results showed that the SVTR has an overall accuracy of 78% and a precision of 92%, performing better than rainfall-based indices in terms of the detection of agricultural drought occurrence. Time-series analysis also showed that SVTR fairly follow the upward and downward trends of rice and corn yield anomalies from the first quarter of 2015 to fourth quarter of 2016. Aside from agricultural drought assessment, the satellite-derived SVTR could be used as an input to drought forecasting models, which may eventually help in decision-making process of crop managers, policymakers, and government institutions toward drought preparedness and mitigation and management.